

# FERRIPROX<sup>®</sup> (deferiprone) TREATMENT TRACKER



Partners in Patient Care



## 12-month treatment tracker (part 1 of 3)

	Baseline	Month 1				Month 2				Month 3				Month 4			
		Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4
FERRIPROX <sup>®</sup> (deferiprone) dose (mg/kg/day)																	
Hemoglobin																	
White blood cell count																	
Neutrophil count																	
Platelet count																	
Medications																	
Serum ferritin																	
Other iron measures*																	
Transfusions (Circle the number of bags of red blood cells you receive)																	

\*How your iron stores will be measured will depend on your healthcare provider.

Please see accompanying Indication and Important Safety Information.

# FERRIPROX<sup>®</sup> (deferiprone) TREATMENT TRACKER



Partners in Patient Care



## 12-month treatment tracker (part 2 of 3)

	Baseline	Month 5				Month 6				Month 7				Month 8			
		Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4
FERRIPROX <sup>®</sup> (deferiprone) dose (mg/kg/day)																	
Hemoglobin																	
White blood cell count																	
Neutrophil count																	
Platelet count																	
Medications																	
Serum ferritin																	
Other iron measures*																	
Transfusions (Circle the number of bags of red blood cells you receive)																	

\*How your iron stores will be measured will depend on your healthcare provider.

Please see accompanying Indication and Important Safety Information.

# FERRIPROX® (deferiprone) TREATMENT TRACKER



Partners in Patient Care



## 12-month treatment tracker (part 3 of 3)

	Baseline	Month 9				Month 10				Month 11				Month 12			
		Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4
FERRIPROX® (deferiprone) dose (mg/kg/day)																	
Hemoglobin																	
White blood cell count																	
Neutrophil count																	
Platelet count																	
Medications																	
Serum ferritin																	
Other iron measures*																	
Transfusions (Circle the number of bags of red blood cells you receive)																	

\*How your iron stores will be measured will depend on your healthcare provider.

Please see accompanying Indication and Important Safety Information.

# FERRIPROX<sup>®</sup> Consumer Indication and Important Safety Information



Partners in Patient Care

## What is FERRIPROX<sup>®</sup> (deferiprone)?

FERRIPROX is a prescription medicine used to treat people with thalassemia syndromes who have iron overload from blood transfusions, when current iron removal (chelation) therapy does not work well enough. FERRIPROX was approved based on its ability to lower iron levels in the blood. There are no studies to show that FERRIPROX improves the symptoms of the disease, or your ability to function, or that it increases the length of time you will live. It is not known if FERRIPROX is safe and effective to treat iron overload due to blood transfusions in people with any other type of anemia that is long lasting, in children, or in people with severe liver problems.

## Important Safety Information

### What is the most important information I should know about FERRIPROX?

FERRIPROX can cause serious side effects, including a very low white blood cell count. One type of white blood cell, called a neutrophil, is important for fighting infections. If you have a low neutrophil count (called neutropenia) you may be at risk of developing a serious infection that can lead to death. Neutropenia is common with FERRIPROX and can become severe in some patients. Severe neutropenia is known as agranulocytosis and if you develop agranulocytosis, you will be at risk of a serious infection that can lead to death.

Your healthcare provider should do a blood test before you start FERRIPROX and weekly during treatment to check your neutrophil count. If you develop neutropenia, your healthcare provider should check your blood counts every day until your white blood count improves. Stop taking FERRIPROX and get medical help right away if you develop symptoms of an infection such as, fever, sore throat or mouth sores, flu-like symptoms, or chills and severe shaking.

### Who should not take FERRIPROX?

Do not take FERRIPROX if you are allergic to deferiprone or any other ingredients in FERRIPROX.

### What should I tell my healthcare provider before taking FERRIPROX?

Tell your healthcare provider if you have liver problems, are pregnant or plan to become pregnant or have any other medical conditions. FERRIPROX can harm your unborn baby so avoid getting pregnant while on therapy. It is not known if FERRIPROX passes into breast milk. You and your healthcare provider should decide if you will take FERRIPROX or breastfeed since you should not do both.

Before taking FERRIPROX tell your healthcare provider if you take other medicines that can lower your neutrophil count or antacids or mineral supplements that contain iron, aluminum, and zinc. Allow at least 4 hours between taking FERRIPROX and products that contain iron, aluminum, or zinc.

### What are other possible side effects of FERRIPROX?

FERRIPROX can cause other serious side effects including increased liver enzyme levels in your blood. Your healthcare provider should do a monthly blood test to check your liver function during treatment.

The most common side effects of FERRIPROX are: reddish brown colored urine, nausea, vomiting, stomach-area (abdominal) pain, joint pain, and a low neutrophil count.

Please see Full Prescribing Information, including Important WARNING and Medication Guide available [here](#) and on [www.ferriproxtotalcare.com](http://www.ferriproxtotalcare.com).